

Cosmic Structure Traced by Precision Measurements of the X-Ray Brightest Galaxy Clusters in the Sky

Thomas H. Reiprich^{*†}, Craig L. Sarazin^{*}, Joshua C. Kempner^{**}, Michael F. Skrutskie^{*}, Gregory R. Sivakoff^{*}, Hans Böhringer[‡] and Jörg Retzlaff[‡]

^{*}*Department of Astronomy, University of Virginia, PO Box 3818, Charlottesville, VA 22903-0818*
[†]*thomas@reiprich.net, <http://www.reiprich.net>*

^{**}*Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, MS-67 Cambridge, MA 02138*

[‡]*Max-Planck-Institut für extraterrestrische Physik, PO Box 1312, 85741 Garching, Germany*

Abstract. The current status of our efforts to trace cosmic structure with 10^6 galaxies (2MASS), 10^3 galaxy clusters (NORAS II cluster survey), and precision measurements for 10^2 galaxy clusters (*HIFLUGCS*) is given. The latter is illustrated in more detail with results on the gas temperature and metal abundance structure for 10^0 cluster (A1644) obtained with XMM-Newton.

Galaxy clusters have been very important tools to study cosmic structure and determine cosmological parameters. Even moderately sized samples yield competitive statistical constraints, e.g., *HIFLUGCS* (Fig. 1), [1]. However, accuracy is currently limited by systematic uncertainties. Luckily major improvements are now feasible, e.g., by taking advantage of new multiwavelength data and especially Chandra and XMM-Newton X-ray observations. We've started a project to tackle systematic uncertainties in flux, temperature, gas and gravitational mass estimates by detailed analyses of the 63 X-ray brightest galaxy clusters in the sky (*HIFLUGCS*). Many of these clusters have already been observed by Chandra and XMM-Newton. Data are accumulating in the archives. The first out of further nine approved Chandra observations proposed by us is being carried out today (2002-12-16).

The NORAS II cluster survey contains about 800 X-ray selected galaxy clusters (Böhringer, Retzlaff et al., in prep.). In order to test for systematic effects caused by different selection techniques — which must be well understood to obtain precise cosmological constraints — we started correlating the NORAS II clusters with the 2MASS extended source catalog (the latter shown in Fig. 1, [2]) and color selected point source catalog. Furthermore galaxy overdensities at cluster positions will be determined in order to estimate richness–mass relations and mass-to-light ratios.

In the following we illustrate possible improvements from the X-ray side on an especially tough example. XMM-Newton observations of A1644 show a very complicated surface brightness distribution on all scales (Reiprich et al., in prep.). A main clump and a smaller sub clump are easily identified. The emission surrounding both core regions is highly non-spherical. And the core region of the main clump itself contains a displaced core-within-a-core. How much do X-ray flux, gas, and gravitational mass estimates based on precision measurements differ from simple estimates where, e.g., the




FIGURE 3. Gas temperature (left) and metal abundance (right) estimates of selected regions based on simultaneous spectral fits to background and vignetting corrected spectra from the MOS1, MOS2, and pn detectors aboard XMM-Newton. Note that R21=R27+R30, R23=R28+R31, R25=R29+32, and R19=full annulus–R18; see Fig. 2. Statistical error bars show the 90% confidence level for one interesting parameter.

clump (a radio source) and the movement through intracluster gas of the main clump. Should more detailed modeling confirm the significance of the abundance excess then other possibilities like cooling of intracluster gas onto the moving sub clump (as seen in A1795 on much smaller scales, [4]) would be more difficult to reconcile with the data.

In summary the temperature structure of A1644 shows that this system is quite complicated as indicated by the surface brightness structure. Interestingly no evidence for substructure has been found in optical and near-infrared observations [5]. The next step is to attempt an improved mass estimate and compare it to more simple estimates as generally applied to clusters at higher redshift and in larger samples. Note that this detailed study required only 12 ks of good data. Chandra and XMM-Newton observations of the complete *HIFLUGCS* are quite cheap but offer a great opportunity for clusters in the era of precision cosmology. Comparison of cluster X-ray data to the wealth of new multiwavelength data, e.g., 2MASS, will also help to reduce systematic uncertainties.

We acknowledge the benefit from the dedicated work of the NORAS II and 2MASS teams. This work was supported by NASA XMM-Newton Grant NAG5-10075. THR acknowledges support by the Celerity Foundation through a Post-Doctoral Fellowship. The XMM-Newton project is an ESA Science Mission with instruments and contributions directly funded by ESA Member States and the USA (NASA). This publication makes use of data products from 2MASS, which is a joint project of the University of Massachusetts and IPAC/Caltech, funded by NASA and NSF.

REFERENCES

1. Reiprich, T. H., and Böhringer, H., *ApJ*, **567**, 716–740 (2002).
2. Jarrett, T. H., Chester, T., Cutri, R., Schneider, S., Skrutskie, M., and Huchra, J. P., *AJ*, **119**, 2498–2531 (2000).
3. Finoguenov, A., Reiprich, T. H., and Böhringer, H., *A&A*, **368**, 749–759 (2001).
4. Fabian, A. C., Sanders, J. S., Ettori, S., Taylor, G. B., Allen, S. W., Crawford, C. S., Iwasawa, K., and Johnstone, R. M., *MNRAS*, **321**, L33–L36 (2001).
5. Tustin, A. W., Geller, M. J., Kenyon, S. J., and Diaferio, A., *AJ*, **122**, 1289–1297 (2001).